



# **Surgical site infection surveillance protocol**

**Edition 7.1**

**(Updated May 2019)**



**Appendix 7-  
Setting up  
surveillance**

## Contents

Setting up surveillance .....	3
1.1 Introduction to SSI Surveillance .....	3
1.2 Who should be involved? .....	4
1.2 a Key Roles.....	4
1.3 Points to consider before setting up surveillance .....	5
1.3a Flowchart- Setting up surveillance .....	6
1.3b Flowchart- National data collection process .....	7
1.3c Flowchart- Patient pathways for standard SSI surveillance.....	8
1.3 d Flowchart – Patient pathways for Light SSI surveillance.....	9
1.4 Data Management .....	10
1.5 Setting up SSI Post-Discharge Surveillance .....	11
1.5a Introduction .....	11
1.5b Requirements for post discharge SSI surveillance .....	11
1.5c Flowchart- Model for all procedures except for caesarean section.....	12
1.5d Flowchart- The caesarean section model.....	13
1.5e Implications for practice.....	14
References.....	15

# Setting up surveillance

## 1.1 Introduction to SSI Surveillance

Infections acquired in hospital are recognised as being associated with significant morbidity. They result in increased length of hospital stay, pain, discomfort and sometimes prolonged or permanent disability, and on occasions result in death.<sup>1,2</sup>

The prevalence of Health Associated Infection (HAI) in 2011 within acute hospitals in Scotland was found to be 4.9%. Urinary tract infection (UTI) was the most prevalent type of HAI (22.6% of all HAI) with Surgical Site Infection (SSI) making up a large proportion of HAI (18.6%), the procedures of particular note were orthopaedic, vascular, and gastrointestinal surgery.<sup>3</sup> The development of HAI are estimated to double the length of post operative stay in hospital significantly increase the cost of care<sup>2,4</sup>.

A surveillance programme is required to establish the cumulative incidence rates of SSI, both nationally and locally. The key aim of SSI surveillance is to improve the quality of patient care and a national surveillance programme is able to provide participating hospitals with robust rates in order for them to compare with similar hospitals and against benchmark rates. Benchmarks of SSI can be a driver for effecting change but require effort and coordination to develop.<sup>5</sup> Evidence demonstrates reductions in SSI rates in hospitals participating in benchmarking projects.<sup>6-8</sup> The evidence also suggests that actively feeding back data to clinicians contributes to reductions in rates of infection.<sup>9</sup>

**The aims of an SSI surveillance programme are to:**

**At a local level:**

- To improve the quality of patient care by collecting surveillance data on the chosen procedure to permit estimation of the magnitude of the risks in hospitalised patients (or the population under examination), and use this data to inform clinical practice improvements

- Analyse and report SSI surveillance data and describe trends in infection rates, to enable clinicians to review or change practice locally that may be required to reduce SSI rates
- Provide timely active feedback of SSI rates to assist clinical teams in minimising the occurrence of SSI.

**At a national level:**

- Monitor trends, including the detection of outbreaks, provide early warning and investigation of problems and subsequent planning and intervention to control
- Examine the impact of interventions to determine best practice
- Gain information on the quality of care to drive improvements
- Prioritise the allocation of resources to promote the reduction in the rate of SSI's.

**1.2 Who should be involved?**

Surveillance primarily involves clinical staff, including infection control teams.

***The local SSI surveillance group should include representation of some or all of the following:***

Infection Control Nurses	Infection Control Managers
Infection Control Doctors/microbiologists	Clinical Effectiveness teams
Medical staff, from relevant disciplines	Theatre staff
Clinical Nurse Specialists	Clinical audit staff
Ward nurses	IT staff
Surveillance Nurses	Clerical and secretarial staff
Community staff	

**1.2a Key Roles**

There are two key roles, which ensure effective surveillance:

### ***Key role 1: Local SSI surveillance co-ordinator.***

The local co-ordinator will be a member of the infection control team or another member of staff with strong links with infection control, for example a member of the clinical effectiveness team. Their key functions are anticipated to be strategic at this local level and include:

- Facilitating the surveillance process
- Engaging with the clinical teams to ensure their continued involvement to drive forward SSI reduction in their surgical procedures
- Provide overall co-ordination and liaison with HPS including ensuring mechanisms are in place for data collection, collation, transfer and dissemination
- Provide local training for staff involved in the surveillance process to ensure the consistent application of SSI definitions and data collection
- Active feedback of SSI data to local stakeholders.

### ***Key role 2: Data transfer co-ordinator.***

The data transfer co-ordinator's key functions are to:


- Ensure that any electronic format utilised locally complies with the database specifications
- Ensure data are correctly entered on the web based reporting system.

## **1.3 Points to consider before setting up surveillance**


- The following sub sections of 1.3 are example flowcharts to consider and assist with setting up of surveillance.
- Specifically, the first flow chart represents the steps necessary to prepare for the commencement of SSI surveillance, followed by the national data collection process. Additional flowcharts contain examples of patient

pathways for specific SSI surveillance including SSI post discharge surveillance, whether an SSI occurs or not during any part of the pathway.


### 1.3a Flowchart- Setting up surveillance




### 1.3b Flowchart- National data collection process



1.3c Flowchart- Patient pathways for standard SSI surveillance



1.3d Flowchart – Patient pathways for Light SSI surveillance



**NB: If at any point of surveillance an SSI is confirmed, apply criteria as per standard surveillance methodology.**

## 1.4 Data Management

- Consideration should be given as to how data will be managed at the setting up stage of programmes.
- Data collection tools can be devised and managed locally, in conjunction with the SSHAIP team at HPS.
- There are three ways in which data can be collected and acquired:
  - The use of automated form processing locally.
  - Web Based reporting.

## **1.5 Setting up SSI Post-Discharge Surveillance**

### **1.5a Introduction**

Due to advances in surgical techniques, length of stay following surgery has decreased.<sup>10</sup> As the length of post operative in patient stay affects the numbers of infections detected, it is recommended that post discharge surveillance (PDS) is carried out, in particular for, those procedures that have an expected short length of hospital stay. The Scottish Government took cognisance of this and since April 2009 it has been mandatory to conduct post discharge SSI surveillance for caesarean section to 10 days post procedures.<sup>11</sup> For hip arthroplasty procedures it is required to conduct readmission surveillance to 30 post operative days.<sup>12</sup> More detail can be found in Mandatory requirements of SSI surveillance <https://www.hps.scot.nhs.uk/web-resources-container/surgical-site-infection-surveillance-protocol-and-resource-pack-edition-71/>

### **1.5 b Requirements for post discharge SSI surveillance**


If valid and reliable data on post discharge SSI's are to be obtained, and if the information collected by different centres is to be comparable, the methods adopted should have the same degree of completeness of coverage of the population at risk, should identify these infections using the same definition and have similar levels of diagnostic accuracy. The gold standard for PDS being that the wound is observed by staff trained in standard SSI surveillance definitions.

For national mandatory (caesarean section) PDS standardised observational methods for identifying SSI and for diagnosis should be in place in all hospitals.


The mandatory requirements in Scotland are:

- Readmission surveillance up to 30 days post-operatively for all procedure except caesarean section (see flowchart 1.5c).
- PDS utilising the normal patient pathway to day 10 post-operatively for all planned and unplanned caesarean section procedures performed (see flowchart 1.5d).

### 1.5c Flowchart- Model for all procedures except for caesarean section



### 1.5d Flowchart- The caesarean section model



### **1.5 e Implications for practice**

- To ensure comparable validity of data the same method i.e., direct observation, for identifying and diagnosing SSI should be adopted at all hospitals undertaking readmission and PDS.
- The SSI definitions applied before and after discharge are identical.
- Those monitoring the surgical wounds must be trained in the monitoring, and diagnosing of SSI according to the standard definitions.
- The duration of surveillance should be the same at all hospitals undertaking readmission and PDS of SSI.

## References

1. Coello R, Charlett A, Wilson J, Ward V, Pearson A, Borriello P. Adverse impact of surgical site infections in English hospitals. *J Hosp Infect* 2005;60:93-103.
2. Plowman R, Graves N, Griffin MA, et al. The rate and cost of hospital-acquired infections occurring in patients admitted to selected specialties of a district general hospital in England and the national burden imposed. *J Hosp Infect* 2001;47:198-209.
3. Health Protection Scotland. Scottish National Point Prevalence Survey of Healthcare Associated Infection and Antimicrobial Prescribing 2016. <https://www.hps.scot.nhs.uk/web-resources-container/national-point-prevalence-survey-of-healthcare-associated-infection-and-antimicrobial-prescribing-2016/> (2017).
4. Broex EC, van Asselt AD, Bruggeman CA, van Tiel FH. Surgical site infections: how high are the costs? *J Hosp Infect* 2009;72:193-201.
5. Gaynes RP. Surgical-site infections (SSI) and the NNIS Basic SSI Risk Index, part II: room for improvement. *Infect Control Hosp Epidemiol* 2001;22:266-7.
6. Gastmeier P, Sohr D, Schwab F, et al. Ten years of KISS: the most important requirements for success. *J Hosp Infect* 2008;70 Suppl 1:11-6.
7. Geubbels EL, Bakker HG, Houtman P, et al. Promoting quality through surveillance of surgical site infections: five prevention success stories. *Am J Infect Control* 2004;32:424-30.
8. Rioux C, Grandbastien B, Astagneau P. Impact of a six-year control programme on surgical site infections in France: results of the INCISO surveillance. *J Hosp Infect* 2007;66:217-23.
9. Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR. Guideline for Prevention of Surgical Site Infection, 1999. Centers for Disease Control and Prevention (CDC) Hospital Infection Control Practices Advisory Committee. *Am J Infect Control* 1999;27:97-132.
10. Ward VP, Charlett A, Fagan J, Crawshaw SC. Enhanced surgical site infection surveillance following caesarean section: experience of a multicentre collaborative post-discharge system. *J Hosp Infect* 2008;70:166-73.

11. A revised framework for national surveillance of healthcare associated infection and the introduction of a new health efficiency and access to treatment (HEAT) target for *Clostridium difficile* Associated Disease (CDAD) for NHS Scotland.CEL(2009)11. Edinburgh: SEHD.\_  
<https://www.sehd.scot.nhs.uk/details.asp?PublicationID=2913>. 2009.
12. A revised framework for national surveillance of healthcare associated infection in Scotland.HDL(2006)38:SEHD;2006. SEHD.\_  
<https://www.sehd.scot.nhs.uk/details.asp?PublicationID=1968>. 2006.